

# Non-arginine-aspartate (non-RD) kinases are associated with innate immune receptors that recognize conserved microbial signatures

Chris Dardick<sup>1</sup>, Benjamin Schwessinger<sup>2</sup> and Pamela Ronald<sup>2</sup>

An important question in the field of plant–pathogen interactions is how the detection of pathogens is converted into an effective immune response. In recent years, substantial insight has been gained into the identities of both the plant receptors and the microbial molecules they recognize. Likewise, many of the downstream signaling proteins and transcription factors that activate defense responses have been characterized. However, the early molecular events that comprise ‘recognition’ and how defense signaling specificity is achieved are not as well understood. In this review we discuss the significance of non-arginine-aspartate (non-RD) kinases, a subclass of kinases that are often found in association with pattern recognition receptors (PRRs).

## Addresses

<sup>1</sup> USDA-ARS, Appalachian Fruit Research Station, 2217 Wiltshire Rd., Kearneysville, WV 25430, United States

<sup>2</sup> Department of Plant Pathology and the Genome Center, University of California, 1 Shields Ave., Davis, CA 95616, United States

Corresponding author: Dardick, Chris ([chris.dardick@ars.usda.gov](mailto:chris.dardick@ars.usda.gov))

Current Opinion in Plant Biology 2012, 15:358–366

This review comes from a themed issue on **Biotic interactions**

Edited by **Pamela C Ronald** and **Ken Shirasu**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 31st May 2012

1369-5266/\$ – see front matter, Published by Elsevier Ltd.

<http://dx.doi.org/10.1016/j.pbi.2012.05.002>

## Introduction

Plant and animal innate immune systems depend on a diverse assortment of cell surface and cytoplasmic receptors that detect and respond to invading pathogens. In plants, these receptors are commonly classified into a group that recognizes conserved microbial signatures (called pattern recognition receptors, PRRs) and a group that recognizes highly variable effectors (nucleotide binding site-leucine rich repeat receptors, NBS-LRRs) [1–3]. The first group contains both extracellular membrane bound receptors and intracellular receptors that frequently contain (or associate with) non-arginine-aspartate (non-RD) kinases. The second group includes intracellular NBS-LRR receptors, which are often fused to additional domains but typically lack kinase domains. These two systems of microbial

perception in plants are commonly referred to as PTI (Pathogen Associated Molecular Pattern [PAMP] Triggered Immunity) and ETI (Effector Triggered Immunity), respectively [1]. Understanding how microbial signals are converted into PTI or ETI remains a fundamentally important issue.

Most plant and animal PRRs identified to date either contain kinase domains or associate with kinases via adaptor molecules [4]. From a simplistic viewpoint, kinases serve as switches that are turned on or off via conformational changes induced by ligand binding. One critical regulatory feature of kinases is the activation loop, which becomes phosphorylated and structurally reoriented to enable substrate access and/or to enhance phosphotransfer efficiency [5]. We previously found that most PRR kinases or PRR-associated kinases contain a change in a conserved arginine (R) located adjacent to the key catalytic aspartate (D) (the so-called RD motif) that facilitates phosphotransfer [6]. This positively charged R residue typically lies within a charge cluster that inhibits catalysis by the neighboring negatively charged D residue. This inhibition can be removed by phosphorylation of the kinase activation loop that lies in close proximity to the RD motif (between kinase subdomains VII and VIII). Activation loop phosphorylation produces negatively charged phospho-amino acids that neutralize the positively charged R residue resulting in kinase activation [6]. Plant and animal kinases associated with recognition of conserved microbial signatures lack the R, in its place having an uncharged residue such as Cys, Gly, Phe, or Leu. Such kinases are referred to as non-RD [7]. The functional significance of non-RD motifs and whether or not charge neutralization plays a role in the catalytic activity of non-RD kinases is currently unknown.

To date, approximately 75 plant receptor-like kinases (RLKs) have been functionally characterized. Nearly one dozen of these are non-RD kinases, all of which have known or putative functions in the recognition of conserved microbial signatures characteristic of PRRs (Figure 1). Unlike their more common RD counterparts, non-RD kinases do not generally auto-phosphorylate the activation loop; presenting a potential mechanistic difference in their activation and/or function [7]. Such changes to the RD motif are likely adaptive changes that may reflect distinctive properties of PRR-mediated signaling.

Figure 1

Kinase	Class	Subfamily	Plant	Pathogen		Reference(s)
LRK10	non-RD	LRK10L-2	Wheat	Fungal		[71]
PR5K	non-RD	LRK10L-2	Arabidopsis	?		[72]
TaRLK 1,2,3	non-RD	LRK10L-2	Wheat	Fungal		[73]
BSR1	non-RD	LRK10L-2	Rice	Fungal/bacterial		[74]
XA21	non-RD	LRR XII	Rice	Bacterial		[14]
XA26	non-RD	LRR XII	Rice	Bacterial		[75]
FLS2	non-RD	LRR XII	Arabidopsis	Bacterial		[15]
EFR	non-RD	LRR XII	Arabidopsis	Bacterial		[16]
DS1	non-RD	LRRXII	Sorghum	Fungal		[76]
CERK1	RD	LysM-I	Arabidopsis	Fungal/bacterial		[44]
RPG1	non-RD	RLCK-OS2	Barley	Fungal		[25]
Pi-d2	non-RD	SD-2b	Rice	Fungal		[77]
LecRK1	non-RD	SD-2b	tobacco	Hornworm		[78]
NgRLK1	non-RD	SD-2b	tobacco	Fungal		[79]
WKS1	non-RD	WAKL-OS	Rice	Fungal		[29]
<b>RLP class PRRs</b>						
CEBiP	none	LysM RLP	Rice	Fungal		[49]
LYM1/3	none	LysM RLP	Arabidopsis	Bacterial		[47]
Ve1	none	LRR RLP	Tomato	Fungal		[38]
LeEIX1/2	none	LRR RLP	Tomato	Yeast		[39]
<b>DAMP Receptors</b>						
WAK1	RD	WAK	Arabidopsis	Oligogalacturonides		[57]
PEPR1	RD	LRR-XI	Arabidopsis	Plant peptide		[54,55]
THESEUS	RD	CrRLK1L-1	Arabidopsis	?		[56]
<b>ETI Receptors</b>						
RPG5	Non-RD	NBS-LRR	Wheat	Fungal		[64]

**Kinase domains**

- Non-RD
- RD
- Non-functional

**Receptor domains**

- Thaumatin/cys rich
- LRR
- LysM
- B-Lectin
- S-locus glycoprotein
- PAN/Apple
- Start
- NBS

Current Opinion in Plant Biology


Table of known and putative PRR receptors along with their domain organizations and kinase functionality (when applicable) [71–79].

## Evolution of plant PRR kinases

The kinase domains found in plant PRRs belong to the same family as those used by their animal counterparts. This kinase family has been dubbed the IRAK/Pelle family in animals (after the human and fly members) and the RLK/Pelle family in plants [8,9]. Animals have very few IRAK/Pelle kinases. Humans have four and flies have just one [10]. Nevertheless, they are still central to both innate immune signaling and development. But unlike plants, in animals, the IRAK/Pelle kinases are all cytoplasmic and lack receptor domains. For example, the cytoplasmic non-RD kinase IRAK1 physically associates with both membrane bound and cytoplasmic PRRs via adaptor proteins to transduce defense signals. A series of phosphorylation events trigger the formation of secondary IRAK1 signaling complexes that dissociate from the receptor complex and ultimately lead to activation of the transcription factor NF $\kappa$ B and mobilization of the innate immune response [11,12<sup>•</sup>].

The RLK/Pelle family in plants underwent a dramatic expansion during early in plant evolution. Concurrent with this expansion, the IRAK domain was fused to numerous extracellular receptor and signaling domains to produce an extremely large repertoire of genes [9,13<sup>••</sup>]. The *Arabidopsis* genome encodes over 600 proteins that contain IRAK kinase domains (76% RLKs) while the rice genome encodes over 1000 (83% RLKs) [4,9]. Of these, IRAK kinases with non-RD motifs make up a minority portion [47 in *Arabidopsis* (8%) and 371 in rice (35%)] yet are abundant in a handful of subfamilies that contain many of the known PRRs. Collectively these PRR subfamilies do not form a distinct clade and are instead scattered among numerous receptor and cytoplasmic kinase clades involved in a wide range of biological functions (Figure 2). Thus, the phylogenetic history of these genes suggests that PRR function probably evolved multiple times. Kinases that function as PRRs or in association with PRRs are nearly indistinguishable from those that control numerous other biological processes, both with respect to their structure and their evolutionary history. To date, the only distinguishing feature that has been identified is the non-RD motif [4]. This implies that there is the potential for kinases that normally function in other processes to adopt PRR functions. It stands to reason that upon adopting a PRR function, additional fine-tuning would be required either through the swapping of kinase domains via recombination or the accumulation of mutations in residues associated with kinase function and/or protein–protein interactions. The finding that many PRRs have adopted changes in the otherwise highly conserved RD kinase motif is probably a consequence of such adaptations. This strict association between the presence of the non-RD motif in kinases associated with PRRs suggests that PRR signaling may have unique requirements that differ from other biological surveillance programs.

Figure 2



PRR RLK subfamilies carrying non-RD kinase domains do not form a unified clade and are scattered among numerous RLK clades involved in various biological functions. Depicted is a reconstructed topological phylogenetic tree from [4] using all known *Arabidopsis* and rice kinase sequences. RLK/Pelle kinase nodes are in black while all non-RLK/Pelle kinase families are shown in gray. Non-RD PRR subfamilies are colored red and labeled. The LysM-I family containing the RD kinase CERK1 is shown in blue.

## Known PRRs contain non-RD kinases

Based on both the presence of non-RD kinase motif and whether or not the subfamilies had undergone lineage specific expansion, we previously predicted that a handful of the >50 RLK subfamilies in rice (ca. 37% of RLKs) and *Arabidopsis* (ca. 7.5% of RLKs) function as PRRs [4]. These include the subfamilies LRR-XII, SD-2a, SD-2b, RLK10L-2 (also known as PR5K), WAK (non-RD subgroups a, c), and WAKL-OS. Since that time, additional confirmed or putative PRRs have been identified for the LRR-XII, SD-2b, RLK10L-2, and WAKL-OS RLK subfamilies (Figure 1). Surprisingly, the largest of these subfamilies including LRR-XII, SD-2b, and RLK10L-2 have remained small in *Arabidopsis* (10, 7, and 13 genes, respectively) relative to rice and poplar that contain >40–100 genes for each of these three subfamilies [4,13<sup>••</sup>]. Both the LRR-XII and SD-2b subfamilies are more ancient and have also remained relatively small in the more primitive bryophyte *Physcomitrella patens* that has 4 LRR-XII and 13 SD-2b kinases [13<sup>••</sup>].

Differences in the expansion rates of RLK subfamilies associated with immune and/or stress responses have been attributed to diversifying selection pressures such as those imposed by pathogens, though substantial direct functional evidence is lacking [4,8,9,13<sup>••</sup>]. The most well

characterized PRR subfamily is LRR XII that includes several known rice and Arabidopsis PRRs including *Xanthomonas* resistance 21 (XA21), Flagellin-Sensing 2 (FLS2), and Elongation Factor-Tu Receptor (EFR) [14–17]. Functional orthologs of Arabidopsis FLS2 have been found in rice, tomato, and tobacco [18–20]. While these orthologs all detect flagellin, their precise specificities vary with regard to the epitope and bacterial strain. The closely related PRR, XA21 recognizes the conserved sulfated protein, Ax21, which is produced by the bacterial pathogens in the genera *Xanthomonas*. Ax21 plays a critical role in bacterial communication (aka ‘quorum sensing’) [21]. Arabidopsis FLS2 recognizes exogenously applied Ax21 peptide derivatives [22]. However, the converse is not true. Rice varieties lacking Xa21 but containing FLS2 do not recognize Ax21. The observation that flagellin and Ax21 perception require distinct receptors in rice suggest that PRRs have diversified in rice as compared with Arabidopsis [3,14,22]. However, it is currently unclear how many different ligands each of these receptors are capable of recognizing and just how specialized they really are. Arabidopsis FLS2 was recently shown to recognize the Clavata 3-peptide (CLV3-p), which is also perceived by the CLV1 and CLV2 RLKs to maintain homeostasis of the shoot apical meristem (SAM). [23<sup>••</sup>]. CLV3-p recognition by FLS2 in the SAM renders the stem cells immune to pathogen infection. This finding suggests that each LRR-XII receptor may be able to recognize more than one ligand. The animal PRR counterparts, called Toll-like receptors (TLRs), also carry extracellular LRR domains and recognize a diverse array of pathogen and host ligands. Structural studies have shown that each of the variable repeats among the extracellular LRRs of TLRs can possess unique ligand binding specificities [24<sup>••</sup>]. Further characterization of plant LRR-XII PRRs should shed light on why PRR clades have expanded in some plants and not others.

### Cytoplasmic non-RD PRRs

While most identified PRRs are membrane bound RLKs, some cytoplasmic non-RD kinases also appear to meet the criteria for PRR functions. For example, barley Rpg1 confers broad-spectrum resistance to stem rust caused by the fungal pathogen *Puccinia graminis* [25]. Rpg1 lacks a canonical receptor domain and predominantly resides in the cytoplasm [26]. It consists of two tandem kinase domains; a catalytically active non-RD kinase domain that is preceded by a catalytically inactive kinase domain. Mutations in either kinase domain abrogate resistance function [27]. Nirmala *et al.*, 2011 identified two conserved *Puccinia* proteins that, when applied in combination, were capable of triggering Rpg1 mediated resistance. Rpg1 was found to interact with both proteins *in vitro* suggesting that despite the lack of an identifiable receptor domain, Rpg1 still functions as a PRR [28]. In another example, wheat Kinase-Start 1 (WKS1) was identified as conferring broad-spectrum resistance in a temperature dependent fashion to the fungal pathogen

*Puccinia striiformis* that causes wheat stripe rust. WKS1 is also a cytoplasmic protein that carries a non-RD kinase domain and a START lipid/sterol binding domain similar to the previously identified Enhanced Disease Resistance 2 (EDR2) protein from Arabidopsis [29]. While the mechanism with which WKS1 confers resistance is still unknown, these examples suggest that PRR function is not limited to detection of extracellular conserved microbial signatures.

### Cooperation between RD and non-RD kinases

In humans, the RD kinase IRAK4 is recruited to PRR receptor complexes upon PAMP perception and is required for activation of the non-RD kinase IRAK1. While not yet confirmed *in vivo*, it appears that IRAK4 directly phosphorylates the IRAK1 activation loop [30]. In *Drosophila*, the IRAK1 counterpart, called Pelle, is likewise critical for TOLL mediated innate immune signaling [31,32]. However, the *Drosophila* counterpart of IRAK4 lacks the kinase domain that was apparently lost at some point during its evolutionary history [33<sup>•</sup>]. Instead, evidence suggests Pelle is not trans-phosphorylated but rather auto-phosphorylated in a concentration dependent manner upon ligand recognition by the associated TOLL receptor [34]. In plants, growing evidence suggests that, like humans, PRR signaling often requires the cooperation of regulatory RD kinases.

A subfamily of RLKs that contain RD kinase domains, called Somatic-Embryogenesis Receptor-Like Kinases (SERKs), have been shown to play critical roles in PRR function. BAK1 (also known as SERK3) is known to physically associate with several PRRs including FLS2, EFR, and XA21 [Chen and Ronald, submitted]. Tomato BAK1 orthologs also associate with tomato receptor-like proteins (RLPs) that function in the immune response including the LeEIX1/2 (tomato ethylene-inducing xylanase) and Ve1 (Verticillium resistance) resistance proteins [35–38,39<sup>•</sup>,40]. BAK1 association with these RLPs appears to be triggered upon pathogen perception. LeEIX1/2 bind a conserved fungal molecule. The association of BAK1/SERK3 with LeEIX2 [39<sup>•</sup>,40]. In the case of Ve1, both BAK1/SERK3 and SERK1 are required for defense signaling. Together with the EIX observations, these results indicate that SERKs can play both positive and negative roles in PRR signaling.

Schwessinger *et al.*, 2011 recently showed that BAK1 kinase activity is required for innate immune signaling by FLS2 and EFR but not for complex formation with either FLS2 or EFR upon exposure to their respective ligands [35,41<sup>••</sup>,42]. As is the case with human IRAK1/IRAK4, it is possible that BAK1 and/or other SERK proteins phosphorylate the activation loops of PRR non-RD kinase domains. In this scenario, phosphorylation of the activation loop upon receptor hetero-dimerization may



serve as a switch for the formation of signaling complexes. A similar model has been proposed for IRAK1 function [43]. Once phosphorylated, the activation loop is re-oriented to expose the ATP binding pocket, which also appears to be required for interaction with downstream signaling proteins. Phosphorylation would thus affect the ability of the non-RD kinase to form complexes required for biological function.

### PRRs with RD kinase domains

While the vast majority of intracellular and extracellular PRRs contain non-RD kinase domains, there is at least one exception that carries an RD kinase. The Arabidopsis Chitin Elicitor Receptor Kinase (CERK1) carries an extracellular plant lysine domain (LysM) and an RD kinase domain [44,45]. AtCERK1 binds fungal chitin, a conserved microbial signature. AtCERK1 is also required for perception of a chemically related compound from bacteria called peptidoglycan (PGN), which is also highly conserved. Unlike the case with chitin, PGN is not directly bound by AtCERK1 [46,47<sup>\*</sup>]. Instead PGN binds two LysM extracellular proteins, LYM1 and LYM3, which lack kinase domains. Chitin perception in rice appears to be more like Arabidopsis PGN perception in that chitin binds the membrane bound receptor Chitin Elicitor Binding Protein (CEBiP) that carries two extracellular LysM domains but lacks a kinase domain [48]. To transduce the immune response upon chitin perception, CEBiP still requires OsCERK1 for signaling [49<sup>\*</sup>].

Thus far it is unclear whether chitin and PGN perception associate with a not-yet-identified non-RD kinase like other characterized PRRs. Closely related RLKs that contain LysM motifs also mediate recognition of Nod factors that are derivatives of chitin oligosaccharides produced by symbiotic rhizobacteria [50,51]. Domain swapping studies between the Nod Factor Receptor NFR1 and CERK1 showed that innate immunity vs. nodulation signaling specificity is conferred by two regions within their kinase domains [52<sup>\*\*</sup>]. The first is a short stretch of amino acids within the kinase activation loop. The second is an adjacent YAQ motif within the  $\alpha$ EF/ $\alpha$ F domain known to regulate the conformation of the kinase activation loop. This finding suggests that the switch between innate immune signaling specificity by CERK1 and nodulation by NFR1 is mediated, at least partly, by changes to the activation loop [52<sup>\*\*</sup>]. Thus, in addition to the RD motif, other residues associated with activation loop function may also be under adaptive selection pressures to specify PRR signaling. This presents the possibility that CERK1 has adopted changes directly to the activation loop in lieu of alterations to the RD motif.

### Relationship between DAMP and PAMP signaling

Like animal innate immune systems, plants also contain receptors that respond to Damage Associated Molecular

Patterns (DAMPs), which are produced as a consequence of pathogen infection and perception [53]. DAMPs function to amplify or reinforce innate immune signaling. Characterized plant DAMP receptors all contain RD kinase domains. These DAMP receptors include the Arabidopsis proteins Wall Associated Kinase (WAK1), Pep Receptor (PEPR1), and Theseus [54–56,57<sup>\*</sup>] as well as the predicted rice DAMP receptor, WAK25 [58]. The close relationship of DAMP-mediated and PAMP-mediated responses is reflected in the fact that DAMP perception also often requires BAK1/SERK3 and also in the observations that PAMP and DAMP signals trigger overlapping transcriptional responses. In the case of WAK25, down regulation compromises XA21-mediated Immunity, indicating that WAK25 is a positive regulator of this process [58]. Despite these similarities, important differences have been described. For example, DAMP detection by WAK1 triggers a response that is similar to FLS2 but is weaker with regard to the amplitude of defense gene expression, is not as comprehensive, and does not induce key Salicylic Acid (SA) dependent genes such as PR1, which are a hallmark of PRR-mediated signaling [59,60]. Domain swapping experiments between the RD kinase domain of WAK1 and the non-RD kinase domain of EFR demonstrated that activation of MAMP and PAMP pathways are specific to their respective kinase domains [61<sup>\*</sup>]. Rice contains a large subfamily of WAKs containing non-RD kinase domains (WAKa,c) that have not yet been characterized [4]. These non-RD WAKs may present an opportunity to better understand how changes in the RD motif influence MAMP vs. PAMP signaling.

### Conclusions

Accumulated evidence supports our earlier finding that PRRs most often contain or associate with non-RD kinases. However, the reason for this association is still unresolved. A key question is whether the selection pressures that drive the adoption of non-RD motifs in PRR kinases are imposed by pathogens, by defense signaling constraints within the host, or by both. It is known that PRR kinases and those involved in downstream signaling events are common targets of pathogen effector proteins that aim to inhibit PRR-mediated signaling by blocking kinase function. For example, the kinase domains of PRRs such as FLS2 are targeted by bacterial effectors that disrupt immune signaling [62]. Such pressures could potentially lead to a diminished role for kinase function. In fact, the kinase auto-phosphorylation activities of Xa21, FLS2, and EFR are relatively weak compared to their RD counterparts like BAK1 [39,63]. While a potentially attractive hypothesis, it seems unlikely or at least incomplete given that numerous other conserved residues that are absolutely required for kinase catalytic activity remain unchanged in PRRs.

Instead of pathogen imposed pressures, innate immune signaling itself may have unique mechanistic requirements that are not satisfied by the canonical RD kinase motif. Innate immune signaling probably requires exquisite control to prevent inadvertent activation. ETI (and less commonly PTI) are often accompanied by programmed cell death (PCD). This is supported by the finding that most non-RD kinases in humans regulate innate immunity, apoptosis, and/or cell cycle control [4]. But if non-RD kinases are important for innate immune regulation in general, then it would be anticipated they would also function in mediating ETI receptor signaling. Two recent studies suggest they may. The barley fungal resistance protein Rpg5 encodes an NBS-LRR protein that also contains a C-terminal non-RD kinase domain [64<sup>••</sup>]. This finding implies that cytoplasmic non-RD kinases could potentially be involved in mediating ETI receptor signaling (at least in some cases). Likewise, the Arabidopsis ETI receptors RPM1, RPS2, and RPS5 were found to all associate with FLS2, forming a larger signaling complex [65]. Thus non-RD containing PTI receptors and ETI receptors may work in concert to trigger innate immunity.

How do non-RD kinases function differently from their RD counterparts? As stated earlier, one possibility is the lack of auto-phosphorylation of the activation loop, which does not commonly occur in non-RD kinases and signifies a potential mechanistic difference in their mode of activation. Currently, the evidence for a lack of activation loop auto-phosphorylation is meager as there are few studies in which *in vivo* phosphorylation sites for non-RD PRRs have been identified. In the case of Xa21 at least, *in vitro* phosphorylation studies failed to identify phospho-amino acids within the activation loop [63,66]. As appears to be the situation in humans, plant PRR kinases may require a regulatory RD kinase such as BAK1 to promote signaling. These regulatory RD kinases may act analogous to IRAK4 and trans-phosphorylate the activation loops of the PRR non-RD kinases. But the consequences of such a mechanism on signaling are still unclear since charge neutralization of the R residue is presumably unnecessary. A scenario proposed for human IRAK1 is that it predominantly plays a scaffolding role for signaling complex formation as data suggest PRR signaling does not depend on IRAK1 kinase activity [43]. Support for such a model in plant PRRs is mixed. FLS2, EFR, and XA21 do have relatively low kinase auto-phosphorylation activities. Likewise, mutations in key catalytic residues of the XA21 kinase domain appear to have only partial effects on resistance function [67]. However, equivalent experiments with FLS2 and EFR suggest that catalytic function is indeed required for innate immune signaling [41<sup>••</sup>,68]. These mixed findings suggest that there may be variation in PRR signaling mechanisms.

An alternative operational mechanism is that the non-RD kinases of PRRs may be constitutively active. In

animals, some non-RD kinases maintain a constitutively active conformation owing to the lack of need for charge neutralization of the inhibiting R residue [69]. Constitutive activation could potentially promote a more rapid response to ligand recognition and/or a sustained signaling response; both of which are inherent properties of innate immune signaling. Alternatively, non-RD kinases may be regulated in an opposite manner with auto-phosphorylation causing a constitutively inactive state. Support for this hypothesis comes from observations in rice, where auto-phosphorylation of the XA21 kinase is promoted by the ATPase XB24. This auto-phosphorylation maintains XA21 in the inactive state. Only upon ligand binding does XB24 dissociate, which appears to be the first step in XA21 signaling [70]. Determining the precise role of kinases in PRR signaling and their mechanism(s) of activation will require further biochemical and structural studies to identify how non-RD kinase catalytic activity is regulated, what role (if any) non-RD catalytic activity plays in signaling, and the relationship between non-RD kinases and their RD regulatory partners.

## Acknowledgements

We would like to acknowledge all of our other colleagues whose work we were unable to include but has substantially contributed to our understanding of PRR kinases. NIH GM59962 provided support for this project.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Jones JD, Dangl JL: **The plant immune system.** *Nature* 2006, **444**:323-329.
  2. Ronald PC, Beutler B: **Plant and animal sensors of conserved microbial signatures.** *Science* 2010, **330**:1061-1064.
  3. Schwessinger B, Ronald PC: **Plant innate immunity: perception of conserved microbial signatures.** *Annu Rev Plant Biol* 2012, **63**:451-482.
  4. Dardick C, Ronald P: **Plant and animal pathogen recognition receptors signal through non-RD kinases.** *PLoS Pathog* 2006, **2**:e2.
  5. Adams JA: **Activation loop phosphorylation and catalysis in protein kinases: is there functional evidence for the autoinhibitor model?** *Biochemistry* 2003, **42**:601-607.
  6. Johnson LN, Nobel MEM, Owen DJ: **Active and inactive protein kinases: structural basis for regulation.** *Cell* 1996, **85**:149-158.
  7. Krupa A, Preethi G, Srinivasan N: **Structural modes of stabilization of permissive phosphorylation sites in protein kinases: distinct strategies in Ser/Thr and Tyr kinases.** *J Mol Biol* 2004, **339**:1025-1039.
  8. Shiu SH, Bleeker AB: **Receptor-like kinases from Arabidopsis form a monophyletic gene family related to animal receptor kinases.** *Proc Natl Acad Sci USA* 2001, **98**:10763-10768.
  9. Shiu SH, Karlowski WM, Pan R, Tzeng YH, Mayer KF, Li WH: **Comparative analysis of the receptor-like kinase family in Arabidopsis and rice.** *Plant Cell* 2004, **16**:1220-1234.
  10. Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S: **The protein kinase complement of the human genome.** *Science* 2002, **298**:1912-1934.

11. Cao Z, Henzel WJ, Gao X: **IRAK: a kinase associated with the interleukin-1 receptor**. *Science* 1996, **271**:1128-1231.
12. Flannery S, Bowie AG: **The interleukin-1 receptor-associated kinases: critical regulators of innate immune signalling**. *Biochem Pharmacol* 2010, **80**:1981-1991.  
This review summarizes what is known to date about the role of the four IRAK kinases in mammalian innate immunity. The authors highlight the complex roles of these kinases and the contrasting differences in their functionalities.
13. Lehti-Shiu MD, Zou C, Hanada K, Shiu SH: **Evolutionary history of stress regulation of plant receptor-like kinase/pelle genes**. *Plant Physiol* 2009, **150**:12-26.  
This study characterizes the expansion of RLK subfamilies in Arabidopsis, rice, and poplar. The authors also demonstrate that most RLK subfamilies (including some known PRR subfamilies) are present in the ancestral *Psychomitrella patens* (moss) genome.
14. Song WY, Wang GL, Chen LL, Kim HS, Pi LY, Holsten T, Gardner J, Wang B, Zhai WX, Zhu LH *et al.*: **A receptor kinase-like protein encoded by the rice disease resistance gene, Xa21**. *Science* 1995, **270**:1804-1806.
15. Chinchilla D, Bauer Z, Regenass M, Boller T, Felix G: **The Arabidopsis receptor kinase FLS2 binds flg22 and determines the specificity of flagellin perception**. *Plant Cell* 2006, **18**:465-476.
16. Zipfel C, Kunze G, Chinchilla D, Caniard A, Jones JD, Boller T, Felix G: **Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts Agrobacterium-mediated transformation**. *Cell* 2006, **125**:749-760.
17. Gomez-Gomez L, Boller T: **Flagellin perception: a paradigm for innate immunity**. *Trends Plant Sci* 2002, **7**:251-256.
18. Robatzek S, Bittel P, Chinchilla D, Köchner P, Felix G, Shiu SH, Boller T: **Molecular identification and characterization of the tomato flagellin receptor LeFLS2, an orthologue of Arabidopsis FLS2 exhibiting characteristically different perception specificities**. *Plant Mol Biol* 2007, **64**:539-547.
19. Takai R, Isogai A, Takayama S, Che FS: **Analysis of flagellin perception mediated by flg22 receptor OsFLS2 in rice**. *Mol Plant-Microbe Interact* 2008, **21**:1635-1642.
20. Hann DR, Rathjen JP: **Early events in the pathogenicity of Pseudomonas syringae on Nicotiana benthamiana**. *Plant J* 2007, **49**:607-618.
21. Lee SW, Han SW, Sriyanum M, Park CJ, Seo YS, Ronald PC: **A type I-secreted, sulfated peptide triggers XA21-mediated innate immunity**. *Science* 2009, **326**:850-853.
22. Danna CH, Millet YA, Koller T, Han SW, Bent AF, Ronald PC, Ausubel FM: **The Arabidopsis flagellin receptor FLS2 mediates the perception of Xanthomonas Ax21 secreted peptides**. *Proc Natl Acad Sci USA* 2011, **108**:9286-9291.
23. Lee H, Chah OK, Sheen J: **Stem-cell-triggered immunity through CLV3p-FLS2 signaling**. *Nature* 2011, **473**:376-379.  
Plant stem cells in the shoot apical meristem are known to be immune to most diseases and have long been used to cultivate pathogen-free plants. This work shows that a host encoded peptide is produced in stem cells that binds to and activates FLS2 to activate innate immunity.
24. Lu J, Sun PD: **The structure of the TLR5-flagellin complex: a new mode of pathogen detection, conserved receptor dimerization for signaling**. *Sci Signal* 2012, **25**(216):pe11.  
A number of studies have now shown how animal PRRs structurally associate with PAMPs. Here the authors describe the structure of the human flagellin receptor TLR5 and summarize the basis for different modes of PAMP recognition.
25. Brueggeman R, Rostoks N, Kudrna D, Kilian A, Han F, Chen J, Druka A, Steffenson B, Kleinohs A: **The barley stem rust-resistance gene Rpg1 is a novel disease-resistance gene with homology to receptor kinases**. *Proc Natl Acad Sci USA* 2002, **99**:9328-9333.
26. Nirmala J, Brueggeman R, Maier C, Clay C, Rostoks N, Kannangara CG, von Wettstein D, Steffenson BJ, Kleinohs A: **Subcellular localization and functions of the barley stem rust resistance receptor-like serine/threonine-specific protein kinase Rpg1**. *Proc Natl Acad Sci USA* 2006, **103**:7518-7523.
27. Nirmala J, Drader T, Chen X, Steffenson B, Kleinohs A: **Stem rust spores elicit rapid Rpg1 phosphorylation**. *Mol Plant-Microbe Interact* 2010, **23**:1635-1642.
28. Nirmala J, Drader T, Lawrence PK, Yin C, Hulbert S, Steber CM, Steffenson BJ, Szabo LJ, von Wettstein D, Kleinohs A: **Concerted action of two avirulent spore effectors activates Reaction to Puccinia graminis 1 (Rpg1)-mediated cereal stem rust resistance**. *Proc Natl Acad Sci USA* 2011, **108**:14676-14681.
29. Fu D, Uauy C, Distelfeld A, Blechl A, Epstein L, Chen X, Sela H, Fahima T, Dubcovsky J: **A kinase-START gene confers temperature dependent resistance to wheat stripe rust**. *Science* 2009, **323**:1357-1360.
30. Li S, Strelow A, Fontana EJ, Wesche H: **IRAK-4: a novel member of the IRAK family with the properties of an IRAK-kinase**. *Proc Natl Acad Sci USA* 2002, **99**:5567-5572.
31. Belvin MP, Anderson KV: **A conserved signaling pathway: the Drosophila toll-dorsal pathway**. *Annu Rev Cell Dev Biol* 1996, **12**:393-416.
32. Shelton CA, Wasserman SA: **Pelle encodes a protein kinase required to establish dorsoventral polarity in the Drosophila embryo**. *Cell* 1993, **72**:515-525.
33. Towb P, Sun H, Wasserman SA: **Tube is an IRAK-4 homolog in a Toll pathway adapted for development and immunity**. *J Innate Immun* 2009, **1**:309-321.  
The mechanisms of PRR-mediated innate immunity in mammals and flies has substantial similarities, however one significant difference is the lack of IRAK-4 in Drosophila. These researchers show that an adaptor protein in drosophila called Tube is orthologous to human IRAK4 but lacks the kinase domain.
34. Shen B, Manley JL: **Pelle kinase is activated by autophosphorylation during Toll signaling in Drosophila**. *Development* 2002, **129**:1925-1933.
35. Chinchilla D, Zipfel C, Robatzek S, Kemmerling B, Nürnberger T, Jones JD, Felix G, Boller T: **A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence**. *Nature* 2007, **448**:497-500.
36. Heese A, Hann DR, Gimenez-Ibanez S, Jones AM, He K, Li J, Schroeder JI, Peck SC, Rathjen JP: **The receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants**. *Proc Natl Acad Sci USA* 2007, **104**:12217-12222.
37. Roux M, Schwessinger B, Albrecht C, Chinchilla D, Jones A, Holton N, Malinovskiy FG, Tör M, de Vries S, Zipfel C: **The Arabidopsis leucine rich repeat receptor-like kinases BAK1/SERK3 and BKK1/SERK4 are required for innate immunity to hemibiotrophic and biotrophic pathogens**. *Plant Cell* 2011, **23**:2440-2455.
38. Fradin EF, Abd-El-Halim A, Masini L, van den Berg GC, Joosten MH, Thomma BP: **Interfamily transfer of tomato Ve1 mediates Verticillium resistance in Arabidopsis**. *Plant Physiol* 2011, **156**:2255-2265.
39. Bar M, Sharfman M, Avni A: **LeEix1 functions as a decoy receptor to attenuate LeEix2 signaling**. *Plant Signal Behav* 2011, **6**:455-457.  
BAK1 is known to positively regulate signaling by several PRRs, however this work establishes that BAK1 can also function as a negative regulator. In this case the authors show that binding of EIX to BAK1 promotes association with a decoy receptor LeEIX1 resulting in impaired signaling by LeEIX2.
40. Bar M, Sharfman M, Ron M, Avni A: **BAK1 is required for the attenuation of ethylene-inducing xylanase (Eix)-induced defense responses by the decoy receptor LeEix1**. *Plant J* 2010, **63**:791-800.
41. Schwessinger B, Roux M, Kadota Y, Ntoukakis V, Sklenar J, Jones A, Zipfel C: **Phosphorylation-dependent differential regulation of plant growth, cell death, and innate immunity by the regulatory receptor-like kinase BAK1**. *PLoS Genet* 2011, **7**:e1002046.  
This is the first study to directly compare the activities of RD and non-RD kinases in PRR signaling. The researchers establish that the non-RD PRR kinases FLS2 and EFR have relatively weaker activity than their RD regulatory counterpart BAK1. They also demonstrate that FLS2 and

EFR do not phosphorylate BAK1 *in vitro* and that the kinase activities of all three are not required for ligand induced complex formation.

42. Schulze B, Mentzel T, Jehle AK, Mueller K, Beeler S, Boller T, Felix G, Chinchilla D: **Rapid heteromerization and phosphorylation of ligand-activated plant transmembrane receptors and their associated kinase BAK1.** *J Biol Chem* 2010, **285**:9444-9451.
  43. Kollwe C, Mackensen AC, Neumann D, Knop J, Cao P, Li S, Wesche H, Martin MU: **Sequential autophosphorylation steps in the interleukin-1 receptor-associated kinase-1 regulate its availability as an adaptor in interleukin-1 signaling.** *J Biol Chem* 2004, **279**:5227-5236.
  44. Miya A, Albert P, Shinya T, Desaki Y, Ichimura K, Shirasu K, Narusaka Y, Kawakami N, Kaku H, Shibuya N: **CERK1, a LysM receptor kinase, is essential for chitin elicitor signaling in *Arabidopsis*.** *Proc Natl Acad Sci USA* 2007, **104**:19613-19618.
  45. Wan J, Zhang XC, Neece D, Ramonell KM, Clough S, Kim SY, Stacey MG, Stacey G: **A LysM receptor-like kinase plays a critical role in chitin signaling and fungal resistance in *Arabidopsis*.** *Plant Cell* 2008, **20**:471-481.
  46. Petutschnig EK, Jones AM, Serazetdinova L, Lipka U, Lipka V: **The LysM-RLK CERK1 is a major chitin binding protein in *Arabidopsis thaliana* and subject to chitin-induced phosphorylation.** *J Biol Chem* 2010, **285**:28902-28911.
  47. Willmann R, Lajunen HM, Erbs G, Newman MA, Kolb D, Tsuda K, Katagiri F, Fliegmann J, Bono JJ, Cullimore JV *et al.*: ***Arabidopsis* lysin-motif proteins LYM1 LYM3 CERK1 mediate bacterial peptidoglycan sensing and immunity to bacterial infection.** *Proc Natl Acad Sci USA* 2011, **108**:19824-19829.
- In this paper the authors identified two PGN binding RLPs that mediate PGN induced innate immunity via CERK1.
48. Kaku H, Nishizawa Y, Ishii-Minami N, Akimoto-Tomiyama C, Dohmae N, Takio K, Minami E, Shibuya N: **Plant cells recognize chitin fragments for defense signaling through a plasma membrane receptor.** *Proc Natl Acad Sci USA* 2006, **103**:11086-11091.
  49. Shimizu T, Nakano T, Takamizawa D, Desaki Y, Ishii-Minami N, Nishizawa Y, Minami E, Okada K, Yamane H, Kaku H, Shibuya N: **Two LysM receptor molecules, CEBiP and OsCERK1, cooperatively regulate chitin elicitor signaling in rice.** *Plant J* 2010, **64**:204-214.
- This was the first paper to show that chitin recognition in rice requires two receptors, CERK1 and the RLP CEBiP that serves as the PRR.
50. Gough C, Cullimore J: **Lipo-chitooligosaccharide signaling in endosymbiotic plant-microbe interactions.** *Mol Plant Microbe Interact* 2011, **24**:867-878.
  51. Radutoiu S, Madsen LH, Madsen EB, Felle HH, Umehara Y, Grønlund M, Sato S, Nakamura Y, Tabata S, Sandal N, Stougaard J: **Plant recognition of symbiotic bacteria requires two LysM receptor-like kinases.** *Nature* 2003, **425**:585-592.
  52. Nakagawa T, Kaku H, Shimoda Y, Sugiyama A, Shimamura M, Takanashi K, Yazaki K, Aoki T, Shibuya N, Kouchi H: **From defense to symbiosis: limited alterations in the kinase domain of LysM receptor-like kinases are crucial for evolution of legume-Rhizobium symbiosis.** *Plant J* 2011, **65**:169-180.
- These authors establish for the first time the domains responsible for signaling specificity between the LysM receptors NFR1 and CERK1 that mediate nodulation and innate immunity, respectively. Through domain swaps they show that a region of the activation loop and an adjacent regulatory domain thought to be associated with activation loop conformation are responsible for switching between nodulation and innate immune responses.
53. Aslam SN, Erbs G, Morrissey KL, Newman MA, Chinchilla D, Boller T, Molinaro A, Jackson RW, Cooper RM: **Microbe-associated molecular pattern (MAMP) signatures, synergy, size and charge: influences on perception or mobility and host defence responses.** *Mol Plant Pathol* 2009, **10**:375-387.
  54. Yamaguchi Y, Pearce G, Ryan CA: **The cell surface leucine-rich repeat receptor for AtPep1, an endogenous peptide elicitor in *Arabidopsis*, is functional in transgenic tobacco cells.** *Proc Natl Acad Sci USA* 2006, **103**:10104-10109.
  55. Krol E, Mentzel T, Chinchilla D, Boller T, Felix G, Kemmerling B, Postel S, Arens M, Jeworutzki E, Al-Rasheid KA, Becker D, Hedrich R: **Perception of the *Arabidopsis* danger signal peptide 1 involves the pattern recognition receptor AtPEPR1 and its close homologue AtPEPR2.** *J Biol Chem* 2010, **285**:13471-13479.
  56. Hématy K, Sado PE, Van Tuinen A, Rochange S, Desnos T, Balzergue S, Pelletier S, Renou JP, Höfte H: **A receptor-like kinase mediates the response of *Arabidopsis* cells to the inhibition of cellulose synthesis.** *Curr Biol* 2007, **17**:922-931.
  57. Brutus A, Sicilia F, Macone A, Cervone F, De Lorenzo G: **A domain swap approach reveals a role of the plant wall-associated kinase 1 (WAK1) as a receptor of oligogalacturonides.** *Proc Natl Acad Sci USA* 2010, **107**:9452-9457.
- By swapping the kinase domains of the DAMP receptor WAK1 and FLS2 these researchers demonstrate that innate immune vs. DAMP signaling specificity is conferred by the respective kinase domains.
58. Seo YS, Chern M, Bartley LE, Han M, Jung KH, Lee I, Walia H, Richter T, Xu X, Cao P *et al.*: **Towards establishment of a rice stress response interactome.** *PLoS Genet* 2011, **7**:e1002020.
  59. Ranf S, Eschen-Lippold L, Pecher P, Lee J, Scheel D: **Interplay between calcium signalling and early signalling elements during defence responses to microbe- or damage-associated molecular patterns.** *Plant J* 2011, **68**:100-113.
  60. Galletti R, De Lorenzo G, Ferrari S: **Host-derived signals activate plant innate immunity.** *Plant Signal Behav* 2009, **4**:33-34.
  61. Denoux C, Galletti R, Mammarella N, Gopalan S, Werck D, De Lorenzo G, Ferrari S, Ausubel FM, Dewdney J: **Activation of defense response pathways by OGs and Flg22 elicitors in *Arabidopsis* seedlings.** *Mol Plant* 2008, **1**:423-445.
- This work describes detailed time-course expression profiling studies comparing DAMP and PAMP recognition. They establish key differences between the two pathways, most significantly the lack of a strong SA-mediated late response by DAMPs marked by the lack of PR1 induction.
62. Zhang J, Li W, Xiang T, Liu Z, Laluk K, Ding X, Zou Y, Gao M, Zhang X, Chen S *et al.*: **Receptor-like cytoplasmic kinases integrate signaling from multiple plant immune receptors and are targeted by a *Pseudomonas syringae* effector.** *Cell Host Microbe* 2010, **7**:290-301.
  63. Liu GZ, Pi LY, Walker JC, Ronald PC, Song WY: **Biochemical characterization of the kinase domain of the rice disease resistance receptor-like kinase XA21.** *J Biol Chem* 2002, **277**:20264-20269.
  64. Brueggeman R, Druka A, Nirmala J, Cavileer T, Drader T, Rostoks N, Mirolohi A, Bennypaul H, Gill U, Kudrna D *et al.*: **The stem rust resistance gene *Rpg5* encodes a protein with nucleotide-binding-site, leucine-rich, and protein kinase domains.** *Proc Natl Acad Sci USA* 2008, **105**:14970-14975.
- The vast majority of ETI receptors encode NBS-LRRs that lack kinase domains. This work identifies a novel NBS-LRR that contains a C-terminal non-RD kinase domain that apparently arose through a recent domain fusion.
65. Qi Y, Tsuda K, Glazebrook J, Katagiri F: **Physical association of pattern-triggered immunity (PTI) and effector-triggered immunity (ETI) immune receptors in *Arabidopsis*.** *Mol Plant Pathol* 2011, **7**:702-708.
  66. Chen X, Chern M, Canlas PE, Jiang C, Ruan D, Cao P, Ronald PC: **A conserved threonine residue in the juxtamembrane domain of the XA21 pattern recognition receptor is critical for kinase autophosphorylation and XA21-mediated immunity.** *J Biol Chem* 2010, **285**:10454-10463.
  67. Andaya CB, Ronald PC: **A catalytically impaired mutant of the rice Xa21 receptor kinase confers partial resistance to *Xanthomonas oryzae* pv *oryzae*.** *Phys Mol Plant Pathol* 2003, **62**:203-208.
  68. Gómez-Gómez L, Bauer Z, Boller T: **Both the extracellular leucine-rich repeat domain and the kinase activity of FLS2 are required for flagellin binding and signaling in *Arabidopsis*.** *Plant Cell* 2001, **13**:1155-1163.
  69. Johnson LN, Nobel MEM, Owen DJ: **Active and inactive protein kinases: Structural basis for regulation.** *Cell* 1996, **85**:149-158.



70. Chen X, Chern M, Canlas PE, Ruan D, Jiang C, Ronald PC: **An ATPase promotes autophosphorylation of the pattern recognition receptor XA21 and inhibits XA21-mediated immunity.** *Proc Natl Acad Sci USA* 2010, **107**:8029-8034.
71. Feuillet C, Schachermayr G, Keller B: **Molecular cloning of a new receptor-like kinase gene encoded at the Lr10 disease resistance locus of wheat.** *Plant J* 1997, **11**:45-52.
72. Wang X, Zafian P, Choudhary M, Lawton M: **The PR5K receptor protein kinase from *Arabidopsis thaliana* is structurally related to a family of plant defense proteins.** *Proc Natl Acad Sci USA* 1996, **93**:2598-2602.
73. Zhou H, Li S, Deng Z, Wang X, Chen T, Zhang J, Chen S, Ling H, Zhang A, Wang D, Zhang X: **Molecular analysis of three new receptor-like kinase genes from hexaploid wheat and evidence for their participation in the wheat hypersensitive response to stripe rust fungus infection.** *Plant J* 2007, **52**:420-434.
74. Dubouzet JG, Maeda S, Sugano S, Ohtake M, Hayashi N, Ichikawa T, Kondou Y, Kuroda H, Horii Y, Matsui M *et al.*: **Screening for resistance against *Pseudomonas syringae* in rice-FOX *Arabidopsis* lines identified a putative receptor-like cytoplasmic kinase gene that confers resistance to major bacterial and fungal pathogens in *Arabidopsis* and rice.** *Plant Biotechnol J* 2011, **9**:466-485.
75. Sun X, Cao Y, Yang Z, Xu C, Li X, Wang S, Zhang Q: **Xa26, a gene conferring resistance to *Xanthomonas oryzae* pv. *oryzae* in rice, encodes an LRR receptor kinase-like protein.** *Plant J* 2004, **37**:517-527.
76. Kawahigashi H, Kasuga S, Ando T, Kanamori H, Wu J, Yonemaru J, Sazuka T, Matsumoto T: **Positional cloning of *ds1*, the target leaf spot resistance gene against *Bipolaris sorghicola* in sorghum.** *Theor Appl Genet* 2011, **123**:131-142.
77. Chen X, Shang J, Chen D, Lei C, Zou Y, Zhai W, Liu G, Xu J, Ling Z, Cao G *et al.*: **A B-lectin receptor kinase gene conferring rice blast resistance.** *Plant J* 2006, **46**:794-804.
78. Gilardoni PA, Hettenhausen C, Baldwin IT, Bonaventure G: ***Nicotiana attenuata* LECTIN RECEPTOR KINASE1 suppresses the insect-mediated inhibition of induced defense responses during *Manduca sexta* herbivory.** *Plant Cell* 2011, **23**:3512-3532.
79. Kim YT, Oh J, Kim KH, Uhm JY, Lee BM: **Isolation and characterization of NgRLK1, a receptorlike kinase of *Nicotiana glutinosa* that interacts with the elicitor of *Phytophthora capsici*.** *Mol Biol Rep* 2010, **37**:717-727.